



REVIEW ARTICLE

OPEN ACCESS

MAJOR CONSIDERATIONS AND OUTCOMES OF CLINICAL STUDIES ON VITAMIN D DEFICIENCY IN HUMAN GUT MICROBIOTA: A SYSTEMATIC REVIEW

^{1,2}Ana Valéria Garcia Ramirez and ²Durval Ribas Filho

¹Clinic Ana Valeria (CAV)- Clinical of Nutrition and Health Science, Street Antônio José Martins Filho, 300, 10^ofloor, room 103, São José do Rio Preto SP, 15092-230, Brazil

²Associação Brasileira de Nutrologia (ABRAN)/Brazilian Association of Nutrology, Catanduva/SP, Rua Belo Horizonte, 909 - Centro, Catanduva SP Brazil 15801-150

ARTICLE INFO

Article History:

Received 26th February, 2019
Received in revised form
08th March, 2019
Accepted 06th April, 2019
Published online 29th May, 2019

Key Words:

Gut microbiota.
Vitamin D. Metabolism.
Metabolic diseases.

ABSTRACT

Introduction: There are 2.0 billion overweight and obese people in the world, and Brazil ranks fifth in the world ranking, with an estimated 18.0 million people. The microbiota of the healthy gastrointestinal system presents around 800 species of bacteria. The human microbiome has about 3 million genes in the gastrointestinal tract, corresponding to 150 times more than the human genome. Vitamin D appears to interact with the immune system through its action on the regulation and differentiation of cells such as lymphocytes, macrophages, and natural killer cells. In addition, there is evidence that vitamin D interferes in vivo and in vitro production of cytokines. The vitamin D plays a significant role in modulating the immune system in the intestine, it is possible that its deficiency could impair the function of the intestinal barrier favoring the translocation of endotoxins such as lipopolysaccharides (LPS) into blood circulation. **Objective:** The present study aimed to investigate the main correlations and outcomes of clinical studies on vitamin D and gut microbiota. **Methodology:** Following the criteria of literary search with the use of the MeSH Terms that were mentioned in the item below on "Search strategies", a total of 55 clinical studies were recruited that were submitted to the eligibility analysis and, after that, 27 studies were selected. The search strategy was performed in the PubMed, Embase, Ovid and Cochrane Library, Web of Science, ScienceDirect Journals (Elsevier), Scopus (Elsevier) and OneFile (Gale) databases. **Final considerations:** In summary, there is a modulatory effect of vitamin D status on the intestinal immune system could influence the commensal bacterial composition and vice versa. The relatively greater abundance of gram-negative genera, such as Haemophilus and Veillonella, may be facilitated by low intake and/or concentration of vitamin D. Relatively small proportion of beneficial bacteria, such as Coprococcus and Bifidobacterium, could trigger immune response and inflammation, requiring actions as dependent on 25 (OH) D. It is concluded that vitamin D role in the maintenance of immune homeostasis seems to occur in part by interacting with the intestinal microbiota. Further studies with the appropriate design are desirable to address the hypothesis raised in the present study.

Copyright © 2019, Adriana Letícia dos Santos Gorayeb et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Ana Valéria Garcia Ramirez and Durval Ribas Filho. 2019. "Major considerations and outcomes of clinical studies on vitamin d deficiency in human gut microbiota: a systematic review", *International Journal of Development Research*, 09, (05), 27479-27483.

INTRODUCTION

Obesity represents a multifactorial disease that causes serious public health problems (Luthold, 2017). There are 2.0 billion overweight and obese people in the world (OMS, 2018) and Brazil ranks fifth in the world, with an estimated 18.0 million people, tending to reach 70.0 million individuals (IBGE, 2019).

*Corresponding author: Ana Valéria Garcia Ramirez,
Clinic Ana Valeria (CAV)- Clinical of Nutrition and Health Science, Street Antônio José Martins Filho, 300, 10^ofloor, room 103, São José do Rio Preto SP, 15092-230, Brazil

The microbiota of the healthy gastrointestinal system presents around 800 species of bacteria, and thousands of these microorganisms live symbiotically with fungi, archaea, and viruses, characterizing each human being with maximum concentration in the colon (Rinninella, 2019). This composition is organized into at least three enterotypes depending on various traits, including genetic background, immune phenotype, eating habits, etc (Arumugam, 2011). In this sense, the human microbiome has about 3 million genes in the gastrointestinal tract, corresponding to 150 times more than the human genome (Huttenhower, 2012). The combination of bacterial cells and genes with host cells and genes leads to the

design of a "superorganism." This depends on the appropriate interactions between the intestinal microbiota and the host to achieve and maintain health (De Filippo, 2010). Vitamin D appears to interact with the immune system through its action on the regulation and differentiation of cells such as lymphocytes, macrophages and natural killer cells (NK) (Angelakis, 2012). In addition, there is evidence that vitamin D interferes in vivo and in vitro production of cytokines (Wu, 2011). Among the immunomodulatory effects demonstrated to be decreased production of interleukin-2 (IL-2), interferon-gamma (INF- γ) and tumor necrosis factor (TNF), inhibition of IL-6 expression and inhibition of the production and secretion of autoantibodies by B-lymphocytes (Tchernof, 2013). The vitamin D plays a significant role in modulating the immune system in the intestine, it is possible that its deficiency could impair the function of the intestinal barrier favoring the translocation of endotoxins such as lipopolysaccharides (LPS) into the blood circulation (Dewulf, 2013). LPS are known to promote low-grade inflammation, which predisposes the patient to insulin resistance. Numerous circulating biomarkers have been used to assess clinical inflammation and for research purposes (Dewulf, 2013 and Everard, 2013). Certain compositions of the gut microbiota (GM) have been associated with systemic inflammation and metabolic disorders. Particularly, gram-negative bacteria, which contain LPS in their outer layer, have been shown to stimulate immune response and to provoke metabolic endotoxemia, while other genera, such as bifidobacteria, reduce endotoxemia (Everard, 2013). Despite being gram-negative, Akkermansia was found to improve the intestinal barrier function and to induce beneficial metabolic effects (Hemarajata, 2013).


Vitamin D deficiency and the lack of VDR have been associated with intestinal dysbiosis and increased susceptibility to intestinal diseases (Mani, 2013). Few studies have investigated whether vitamin D status contributes to disorders of glucose metabolism by modulating the composition of the GM (Luthold, 2017 and Arumugam, 2011). A better understanding of the underlying mechanisms of cardiometabolic diseases is important in relation to their impact on population mortality rates. Certain compositions of the GM have been associated with systemic inflammation and metabolic disorders. Particularly, gram-negative bacteria, containing LPS in their outer layer, have been shown to stimulate immune response and to provoke metabolic endotoxemia, while other genera, such as Bifidobacteria, reduce endotoxemia (Yun, 2017). Therefore, using a systematic review, the present study aimed to investigate the main correlations and outcomes of clinical studies on vitamin D and gut microbiota.

MATERIALS AND METHODS

Study design: Following the criteria of literary search with the use of the MeSH Terms that were cited in the item below on "Search strategies", a total of 55 clinical studies were recruited that were submitted to the eligibility analysis and, after that, 27 studies were selected, following the rules of systematic review-PRISMA (Transparent reporting of systematic reviews and meta-analyses-<http://www.prisma-statement.org/>).

Search Strategy and Information Sources: The search strategy was performed in PubMed, Embase, Ovid and Cochrane Library, Web of Science, ScienceDirect Journals (Elsevier), Scopus (Elsevier), OneFile (Gale) followed the

following steps: - search for MeSH Terms: *Gut microbiota. Vitamin D. Metabolism. Metabolic diseases*, use of boolean "and" between mesh terms and "or" among historical findings. All references are registered in EndNote.



LITERATURE REVIEW AND DISCUSSION

The human gastrointestinal tract (HGIT) is the most densely populated organism with commensal and symbiotic microorganisms, mostly bacteria, but also fungi, archaea and viruses (Luthold, 2017), harboring ten times more bacteria than the number of cells that make up our organism (OMS, 2018). Individuals present distinct bacterial compositions, partly defined genetically and in another determined by individual and environmental characteristics, such as mode of birth (normal delivery or cesarean section), age and eating habits, which results in a great deal of intra and interindividual variability (IBGE, 2019). One of the ways to evaluate the communities living in the HGIT is by the taxonomic classification that distributes the bacteria in phyla, classes, order, family, genus and species (Rinninella, 2019). It is estimated that, in the GM, there are about a thousand species, distributed in more than 50 different phyla. Metagenomic studies indicate that, in the human microbiota, there are about 3.3 million different genes, 150 times more than the human genome (Arumugam, 2011). The GM of each individual is characterized specifically by groups of bacteria called enterotypes (Huttenhower, 2012). Three enterotypes are characterized by three groups of dominant bacteria such as *Bacteroides* (enterotipo I), *Prevotella* (enterotipo II), or *Ruminococcus* (enterotipo III). Each enterotipo harbors different genera of bacteria. These three enterotypes are not just enumerations of bacteria, they are also specifically reassembled by functions. In fact, an enterotype is a harmonious association of several species of bacteria rather than a systematic addition of bacterial species (De Filippo, 2010).

Approaches On The Vitamin D and Gut Microbiota: The primary source of vitamin D depends on exposure of the skin to sunlight with up to 20.0% coming from ingestion. It is still

controversial as to whether the consumption of foods containing vitamin D has a direct impact on circulating levels (Angelakis, 2007). Vitamin D₂ (ergocalciferol) is found in yeast, mushrooms and some vegetables and vitamin D₃ (cholecalciferol), synthesized in the skin by ultraviolet radiation, is found in foods of animal origin (Wu, 2011). To be biologically active, vitamin D undergoes hydroxylation in the liver mediated by 25-hydroxylase and in the kidney by 1 α -hydroxylase. 1,25-dihydroxyvitamin D is recognized by its specific receptors (VDR) in different cells, primarily in the gut to increase calcium uptake and in the bone to regulate skeletal homeostasis (Tchernof, 2013). Altered metabolic patterns result in metabolic disorders of calcium and phosphorus with vitamin D disorders having been implicated in some diseases (Tchernof, 2013). Vitamin D plays important roles in innate and adaptive immune responses, in the cell cycle and in metabolic processes, evidenced by the reported relationship between its deficiency and the prevalence of immune-mediated disorders, cancer and cardiometabolic diseases (Dewulf, 2013). An inverse correlation between vitamin D concentrations has been described in respect to the prevalence of obesity and type 2 diabetes mellitus (Everard, 2013; Hemarajata, 2013 and Mani, 2013).

VDR play a part in the production of β cells, endothelium, cardiac myocytes and renin suggesting a role for vitamin D in immune-mediated disorders, cancer and cardiometabolic diseases (Hemarajata, 2013; Mani, 2013; Bai, 2017). In addition, there is evidence that vitamin D deficiency increases inflammatory cytokines and reduces insulin sensitivity with both these conditions having been described as pathophysiological links in cardiometabolic diseases (Yun, 2017). Metabolism-induced GMendotoxemia has been associated with increased cardiometabolic risk (Riva, 2017). Vitamin D deficiency could increase the competitive advantage of *Haemophilus* and *Veillonella*, as these pathogens are found to be relatively more abundant in subjects with low compared to high intake of vitamin D (Borgo, 2017). These gram-negative bacteria could explain the higher levels of LPS detected in people who ingest low levels of vitamin D (Angelakis, 2018). On the other hand, a relatively small proportion of bacteria with beneficial effects - such as *Coprococcus* and *Bifidobacterium*- could activate intestinal immune response and induce local inflammation, requiring anti-inflammatory compensation pathways such as those dependent on 25(OH)D (Belizário, 2018). The *Coprococcus* and *Bifidobacterium* maintained inversely associated with 25 (OH) D concentration after adjustments. Both genders were related to adequate homeostasis of BMI and glucose, as well as production of butyrate and anti-inflammatory actions (Tsai, 2019).

It is possible that exposure to a high abundance of those bacteria can attenuate the immune response. Association *Coprococcus* with vitamin D status was described when the vitamin was supplemented for the treatment of immune-mediated diseases (Luthold, 2017), but not in healthy individuals. In normal participants, the discovery of an association of 25 (OH) D with *Coprococcus* abundance reflects intestinal immune homeostasis and a stable inflammatory systemic state. To the best of our knowledge, no study has shown an association of *Bifidobacterium* with vitamin D status (Luthold, 2017). The attenuation or abolition of the association of these genera with three different inflammatory markers suggests that inflammation may partially mediate the

relationship between vitamin and microbiota concentration. This is in accordance with the reported evidence of its anti-inflammatory action (Tsai, 2019), and with a pro-inflammatory state when vitamin D is lacking (Kayser, 2019). VDR knockout mice exhibited chronic inflammation in the gastrointestinal tract, even in the face of a non-pathogenic microbiota disease. Despite the controversies over a cause-and-effect relationship, there is a consensus that this vitamin is an important mediator of intestinal defenses against infectious diseases agents (Swain Ewald, 2018). Inverse associations between inflammatory markers and 25 (OH) D have been reported. These results support the role of vitamin D in maintaining the homeostasis of the immune system; some authors speculate that this occurs, in part, through interactions with the GM, although the study design excludes establishing cause-and-effect relationships (Vatanen, 2019). It has been previously described that vitamin D deficient rats, exposed to a bacterial pathogen, exhibited increased endotoxin translocation and inflammatory cytokine production. There is a significant inverse correlation of 25(OH)D with E-selectin and C-reactive protein concentrations, suggesting that even among healthy individuals, the vitamin D status may cause an anti-inflammatory condition (Bianchi, 2018). As a matter of fact, several studies have reported that 25(OH)D plays a significant role in the immune system, as low grade inflammation is correlated with higher serum levels as well as higher consumption (Moossavi, 2018). In addition, previous studies show that *Akkermansiamuciniphilais* linked to effects on metabolic and inflammatory profiles (Bell, 2018). Using animal models, *A. muciniphila* benefits intestinal permeability, mucosal layer thickness and the metabolism in obesity and type 2 diabetes (Bell, 2018). Previously, *Haemophilus* was associated with IBD and with levels of LPS, and *Veillonella* with increased inflammation cytokines. These gram-negative bacteria have an outer layer of LPS that is less prominent in subgroups of patients with higher intakes of 25(OH)D (Bell, 2018).

Modulation of Gut Microbiota: Microorganisms that colonize the intestine can alter gene expression in intestinal mucosal cells and ultimately alter the function of human gastrointestinal tract (HGIT) (Luthold, 2013). In general, the gut microbiota (GM) is mostly composed of non-pathogenic and health-promoting bacteria but, to a small extent, by potentially pathogenic bacteria (Luthold, 2017 and OMS, 2018). The diet is a factor determining the characteristics of intestinal colonization, especially vitamin D. This highly influences long-term host phenotypes and is not abruptly altered by short-term interventions (Rinninella, 2019). There are important gaps in the knowledge about how feeding and other life habits could alter the composition of the microbiota and how the microbiota modulates positively or negatively the energy balance. Some diets can modify the pattern of intestinal colonization from the beginning of life (Rinninella, 2019; Arumugam, 2011). It is suggested that feeding has a direct effect on the GM, which would ultimately result in changes in biochemical reactions in the intestinal lumen (Huttenhower, 2012 and De Filippo, 2010). However, the definition of this effect is still something to be explored, since most of the existing studies present confounding factors and/or low comparability by using different experimental models, such as different methods of analysis, species of animals studied, lack of standardization of the degrees of body adiposity, the age of the participants and the diet employed (Tchernof, 2013).

Furthermore, since vitamin D plays a role in modulating the immune system in the intestine, it is possible that its deficiency could impair the intestinal barrier function favoring the translocation of endotoxins such as lipopolysaccharides (LPS) in the circulation (Luthold, 2017). The LPS are known as low-grade inflammation, which predisposes insulin resistance. Numerous circulating biomarkers have been used to assess clinical inflammation and investigation (OMS, 2018). In this context, many efforts have been made to understand the link between GM composition and obesity, as well as the role of food ingredients, such as pro and prebiotics, in modulating the GM (Everard, 2013). Studies involving the GM composition of obese individuals are still controversial, making it difficult to treat obesity (Hemarajata, 2013).

Conclusion

In summary, there is a modulatory effect of vitamin D status on the intestinal immune system could influence the commensal bacterial composition and vice versa. The relatively greater abundance of gram-negative genera, such as *Haemophilus* and *Veillonella*, may be facilitated by low intake and/or concentration of vitamin D. Relatively small proportion of beneficial bacteria, such as *Coprococcus* and *Bifidobacterium*, could trigger immune response and inflammation, requiring actions anti-inflammatory as dependent on 25 (OH) D. It is concluded that vitamin D role in maintaining immune homeostasis seems to occur in part by interacting with the intestinal microbiota. Further studies with the appropriate design are desirable to address the hypothesis raised in the present study.

Conflict of interests: There is no conflict of interest between authors.

Acknowledgment: The authors thank the Clinic Ana Valéria Ramirez and also thank ABRAN - Brazilian Association of Nutrology for all learning and optimization of knowledge.

REFERENCES

Angelakis E, Armougom F, Million M, Raoult D. The relationship between gut microbiota and weight gain in humans. *Future Microbiol.*, 7(1):91-109.

Angelakis E, Bachar D, Yasir M, Musso D, Djossou F, Melenotte C, Robert C, Davoust B, Gaborit B, Azhar EI, Bibi F, Dutour A, Raoult D. Comparison of the gut microbiota of obese individuals from different geographic origins. *New Microbes New Infect.* 2018 Nov 22;27:40-47 [doi: 10.1016/j.nmni.2018.11.005. eCollection 2019 Jan].

Arumugam M, Raes J, Pelletier E, Le Paslier D, Yamada T, Mende DR, *et al.* 2011. Enterotypes of the human gut microbiome. *Nature.*, 473(7346):174-80.

Bai J, Hu Y, Bruner DW. Composition of gut microbiota and its association with body mass index and lifestyle factors in a cohort of 7–18 years old children from the American Gut Project. *Pediatr. Obes.* 2018.

Belizário JE, Faintuch J, Garay-Malpartida M. Gut Microbiome Dysbiosis and Immunometabolism: New Frontiers for Treatment of Metabolic Diseases. *Mediators Inflamm.* 2018 Dec 9; 2018: 2037838 [doi: 10.1155/2018/2037838. eCollection 2018].

Bell, V., Ferrão, J., Pimentel, L., Pintado, M., Fernandes, T. One Health, Fermented Foods, and Gut Microbiota.

Foods. 2018 Dec 3;7(12) [pii: E195. doi: 10.3390/foods7120195].

Bianchi F, Duque ALRF, Saad SMI, Sivieri K. Gut microbiome approaches to treat obesity in humans. *ApplMicrobiolBiotechnol.* 2019 Feb;103(3):1081-1094 [doi: 10.1007/s00253-018-9570-8. Epub 2018 Dec 16].

Borgo, F.; Riva, A.; Benetti, A.; Casiraghi, M.C.; Bertelli, S.; Garbossa, S.; Anselmetti, S.; Scarone, S.; Pontiroli, A.E.; Morace, G.; *et al.* Microbiota in anorexia nervosa: The triangle between bacterial species, metabolites and psychological tests. *PLoS ONE* 2017, 12, e0179739.

De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, *et al.* Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *ProcNatlAcadSci USA.* 2010; 107(33):14691-96.

Dewulf EM, Ge Q, Bindels LB, Sohet FM, Cani PD, Brichard SM, *et al.* Evaluation of the relationship between GPR43 and adiposity in human. *NutrMetab.* 2013;10(1):11.

Everard A, Belzer C, Geurts L, Ouwerkerk JP, Druart C, Bindels LB, *et al.* Cross-talk between *Akkermansiamuciniphila* and intestinal epithelium controls diet-induced obesity. *Proc Natl Acad Sci USA.* 2013; 110(22):9066-71.

Hemarajata P, Versalovic J. Effects of probiotics on gut microbiota: mechanisms of intestinal immunomodulation and neuromodulation. *Ther Adv Gastroenterol.* 2013; 6(1):39-51.

Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, Chinwalla AT, *et al.* Structure, function and diversity of the healthy human microbiome. *Nature.* 2012; 486(7402):207-14.

IBGE- Instituto Brasileiro de Geografia e Estatística. Disponível em: < <http://www.ibge.gov.br>>. Acessado em fevereiro de 2019.

Kayser BD, Lhomme M, Prifti E, Da Cunha C, Marquet F, Chain F, Naas I, Pelloux V, Dao MC, Kontush A, Rizkalla SW, Aron-Wisniewsky J, Bermúdez-Humarán LG, Oakley F, Langella P, Clément K, Dugail I. Phosphatidylglycerols are induced by gut dysbiosis and inflammation, and favorably modulate adipose tissue remodeling in obesity. *FASEB J.* 2019 Jan 4:fj201801897R [doi: 10.1096/fj.201801897R. Epub ahead of print].

Luthold RV, Fernandes GR, Franco-de-Moraes AC, Folchetti LG, Ferreira SR. Gut microbiota interactions with the immunomodulatory role of vitamin D in normal individual s. *Metabolism.* 2017 Apr;69:76-86 [doi: 10.1016/j.metabol.2017.01.007. Epub 2017 Jan 13].

Mani V, Hollis JH, Gabler NK. Dietary oil composition differentially modulates intestinal endotoxin transport and postprandial endotoxemia. *NutrMetab.* 2013;10(1):6.

Moossavi S, Bishehsari F. Microbes: possible link between modern lifestyle transition and the rise of metabolic syndrome. *Obes Rev.* 2019 Mar;20(3):407-419 [doi: 10.1111/obr.12784. Epub 2018 Dec 11].

OMS- Organização Mundial de Saúde. Disponível em: <https://www.scbm.org.br/endoscopia-e-obesidade/> Acesso em: fevereiro de 2018.

Rinninella E, Raoul P, Cintoni M, Franceschi F, Miggianno GAD, Gasbarrini A, Mele MC. What is the Healthy Gut Microbiota Composition? A Changing Ecosystem across Age, Environment, Diet, and Diseases. *Microorganisms.* 2019 Jan 10;7(1) [pii: E14. doi: 10.3390/microorganisms7010014].

- Riva A, Borgo F, Lassandro C, Verduci E, Morace G, Borghi E, Berry D. Pediatric obesity is associated with an altered gut microbiota and discordant shifts in Firmicutes populations. *Environ. Microbiol.* 2017, 19, 95–105.
- Swain Ewald HA, Ewald PW. Natural Selection, The Microbiome, and Public Health. *Yale J Biol Med.* 2018 Dec 21;91(4):445-455. eCollection 2018 Dec.
- Tchernof A, Després JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev.* 2013;93(1):359-404.
- Tsai YL, Lin TL, Chang CJ, Wu TR, Lai WF, Lu CC, Lai HC. Probiotics, prebiotics and amelioration of diseases. *J Biomed Sci.* 2019 Jan 4;26(1):3 [doi: 10.1186/s12929-018-0493-6].
- Vatanen T, Plichta DR, Somani J, Münch PC, Arthur TD, Hall AB, Rudolf S, Oakeley EJ, Ke X, Young RA, Haiser HJ, Kolde R, Yassour M, Luopajarvi K, Siljander H, Virtanen SM, Ilonen J, Uibo R, Tillmann V, Mokurov S, Dorshakova N, Porter JA, McHardy AC, Lähdesmäki H, Vlamakis H, Huttenhower C, Knip M, Xavier RJ. Publisher Correction: Genomic variation and strain-specific functional adaptation in the human gut microbiome during early life. *Nat Microbiol.* 2019 Mar;4(3):545 [doi: 10.1038/s41564-019-0393-x].
- Wu GD, Chen J, Hoffmann C, Bittinger K, Chen Y, Sue A, *et al.* Linking long-term dietary patterns with gut microbial enterotypes. 2011; 334(6052):105-8.
- Yun, Y, Kim HN, Kim SE, Heo SG, Chang Y, Ryu S, Shin H, Kim HL. Comparative analysis of gut microbiota associated with body mass index in a large Korean cohort. *BMC Microbiol.* 2017, 17, 151.
